

CSE 547
Chapter 1, problem 20

Solution 2

Chapter 1 problem 20

Solve the general five-parameter recurrence

- $h(1)=\alpha$;
- $h(2n+j) = 4h(n) + \gamma_j n + \beta_j$, for $j=0,1$ and $n \geq 1$
for $n \in \mathbb{N}$

According to the book, we should ideally solve $h(n)$ for 5 parameters, and find the general equation

$$h(n) = a(n)\alpha + b(n)\beta_0 + c(n)\beta_1 + d(n)\gamma_0 + e(n)\gamma_1.$$

Instead, we're going to take a different approach, using binary numbers.

First 5 solutions:

$$h(1) = \alpha$$

$$h(2) = h(2^*1+0) = 4h(1)+n\gamma_0+\beta_0 = 4\alpha+\gamma_0+\beta_0$$

$$h(3) = h(2^*1+1) = 4h(1)+n\gamma_1+\beta_1 = 4\alpha+\gamma_1+\beta_1$$

$$h(4) = h(2^*2+0) = 4h(2)+n\gamma_0+\beta_0 =$$

$$4(4\alpha+\gamma_0+\beta_0)+2\gamma_0+\beta_0 =$$

$$16\alpha+4\gamma_0+4\beta_0+2\gamma_0+\beta_0 = 16\alpha+6\gamma_0+5\beta_0$$

$$h(5) = h(2^*2+1) = 4h(2)+n\gamma_1+\beta_1 =$$

$$4(4\alpha+\gamma_0+\beta_0) + 2\gamma_1+\beta_1 = 16\alpha+4\gamma_0+4\beta_0+$$

$$2\gamma_1+\beta_1$$

There is no obvious pattern from these steps

We can use binary numbers to find the solution.

We can write out the number k in binary expansion:

$$k = (1 b_{m-1} b_{m-2} \dots b_1 b_0)_2$$

We know that each term b_i is 0 or 1, with the leading digit, α , always 1.

$$h(1) = \alpha$$

$$h(2n+j) = 4h(n) + \gamma_j n + \beta_j \quad \text{for } j=0,1$$

$$h(1 b_{m-1} b_{m-2} \dots b_1 b_0)_2 = 4h(1 b_{m-1} b_{m-2} \dots b_1)_2 + \gamma_{b_0} (1 b_{m-1} b_{m-2} \dots b_1)_2 + \beta_{b_0}$$

$$= 4[4h(1 b_{m-1} b_{m-2} \dots b_2)_2 + \gamma_{b_m}(1 b_{m-1} b_{m-2} \dots b_2)_2 + \beta_{b_1}] + \gamma_{b_1}(1 b_{m-1} b_{m-2} \dots b_2)_2 + \beta_{b_1} + \gamma_{b_0}(1 b_{m-1} b_{m-2} \dots b_1)_2 + \beta_{b_0}$$

As we can see, by expanding the formula, we reduce the number of digits inside each binary term

As we expand the formula completely, the equation becomes

$$h(1 b_{m-1} b_{m-2} \dots b_1 b_0)_2 = 4^m \alpha + 4^{m-1} (1)_2 \gamma_{b_{m-1}} + 4^{m-2} (1 b_{m-2})_2 \gamma_{b_{m-2}} \dots + 4^0 (1 \dots b_1)_2 \gamma_{b_0} + 4^{m-1} \beta_{b_{m-1}} + 4^{m-2} \beta_{b_{m-2}} \dots + 4^1 \beta_{b_1} + 4^0 \beta_{b_0}$$

Where $m+1$ is the number of digits in the binary number $k = (1 b_{m-1} b_{m-2} \dots b_1 b_0)_2 = (b_m \dots b_0)_2$.

$$h(1 b_{m-1} b_{m-2} \dots b_1 b_0)_2 = 4^m \alpha + 4^{m-1} (1)_2 \gamma_{b_{m-1}} + 4^{m-2} (1 b_{m-2})_2 \gamma_{b_{m-2}} \dots + 4^0 (1 \dots b_1)_2 \gamma_{b_0} + 4^{m-1} \beta_{b_{m-1}} + 4^{m-2} \beta_{b_{m-2}} \dots + 4^1 \beta_{b_1} + 4^0 \beta_{b_0}$$

We can combine all the γ terms as a summation, which becomes

$$\sum 4^{m-i} (1 \dots b_{m-i+1})_2 \gamma_{b_{m-i}}$$

summation for $i: 1 \leq i \leq m$.

Similarly we can do the same thing to all the β terms, which turns into

$$\sum 4^{m-i} \beta_{b_{m-i}},$$

summation for $i: 1 \leq i \leq m$

$m+1$ is the number of digits of the binary number

$$k = (1 b_{m-1} b_{m-2} \dots b_1 b_0)_2 = (b_m \dots b_0)_2$$

The closed formula becomes

$$h(b_m \dots b_0) = 4^m \alpha + \sum 4^{m-i} (1 \dots b_{m-i+1})_2 \gamma_{b_{m-i}} + \sum 4^{m-i} \beta_{b_{m-i}}$$

where summation is for $i: 1 \leq i \leq m$, $m+1$ is the number of digits in the binary number

$$k = (1 b_{m-1} b_{m-2} \dots b_1 b_0)_2 = (b_m \dots b_0)_2 h(b_m \dots b_0),$$

Lets try $k=5$, which we solved already

$$h(5) = 16\alpha + 4\gamma_0 + 4\beta_0 + 2\gamma_1 + \beta_1$$

In binary form: $h(5) = (101)$

$$h(5) = 4^m \alpha + \sum 4^{m-i} (1 \dots b_{m-i+1})_2 \gamma_{b_{m-i}} + \sum 4^{m-i} \beta_{b_{m-i}}$$

$$h(5) = 4^2 \alpha + 4^1 (1) \gamma_{b_1} + 4^0 (10) \gamma_{b_0} + 4^1 \beta_{b_1} + 4^0 \beta_{b_0}$$

$$h(5) = 16\alpha + 4\gamma_{b_1} + 1(2) \gamma_{b_0} + 4\beta_{b_1} + \beta_{b_0}$$

$$h(5) = 16\alpha + 4\gamma_0 + 2\gamma_1 + 4\beta_0 + \beta_1$$

$$h(1) = 4^0\alpha + 0\gamma_{b_{m-i}} + 0\beta_{b_{m-i}} = \alpha$$

$$h(2) = 4^1\alpha + 4^0(1)\gamma_{b_0} + \beta_{b_0} = 4\alpha + \gamma_0 + \beta_0$$

$$h(3) = 4^1\alpha + 4^0(1)\gamma_{b_0} + 4^0\beta_{b_0} = 4\alpha + \gamma_1 + \beta_1$$

$$\begin{aligned} h(4) &= 4^2\alpha + 4^1(1)\gamma_{b_1} + 4^0(10)\gamma_{b_0} + 4^1\beta_{b_1} + 4^0\beta_{b_0} \\ &= 16\alpha + 4(1)\gamma_0 + (2)\gamma_0 + 4\beta_0 + \beta_0 \\ &= 16\alpha + 6\gamma_0 + 5\beta_0 \end{aligned}$$

Which are the same values we got recursively.