

Persistent Homology

David Gu

Computer Science Department
Stony Brook University

gu@cs.stonybrook.edu

August 10, 2022

Persistent Homology




Figure: Handle detection by finding the handle loops and the tunnel loops.




Figure: Topological Denoise in medical imaging.

Cech Complex




Figure: Čech complex.

Definition (Cech Complex)

Given a set of points $\{x_\alpha\}$ in Euclidean space \mathbb{R}^n , the Cech complex (also known as the nerve), \mathcal{C}_ϵ , is the abstract simplicial complex where a set of $k + 1$ vertices spans a k -simplex whenever the $k + 1$ corresponding closed $\epsilon/2$ -ball neighborhoods have nonempty intersection.

Definition (Vietoris-Rips Complex)

Given a set of points $\{x_\alpha\}$ in Euclidean space \mathbb{R}^n , the Vietoris-Rips complex, \mathcal{R}_ϵ , is the abstract simplicial complex where a set S of $k + 1$ vertices spans a k -simplex whenever the distance between any pair of points in S is at most ϵ .

Cech Complex






Figure: Cech complex.

Cech Complex




$$\alpha = \alpha_3$$




$$\alpha = \alpha_4$$

Figure: Cech complex.

Cech Complex



$$\alpha = \alpha_5$$



persistent diagram

Figure: Cech complex.

Definition (filtration)

A filtration of a simplicial complex \mathbb{K} is a nested sequence of complexes,

$$\emptyset = \mathbb{K}_{-1} \subset \mathbb{K}_0 \subset \mathbb{K}_1 \subset \cdots \subset \mathbb{K}_n = \mathbb{K}.$$

Example

Suppose \mathbb{K} is a simplicial complex, we sort all the simplices in a sequence

$$\sigma_1^0, \sigma_2^0, \dots, \sigma_{n_0}^0, \sigma_1^1, \sigma_2^1, \dots, \sigma_{n_1}^1, \sigma_1^2, \sigma_2^2, \dots, \sigma_{n_2}^2.$$

where σ_i^k is the i -th k -simplex in \mathbb{K} . Then we relabel all the simplices as

$$\sigma^0, \sigma^1, \sigma^2, \dots,$$

We define \mathbb{K}_j as the union of $\sigma^0, \sigma^1, \dots, \sigma^j$.




Figure: Chain, cycle, boundary groups and their images under the boundary operators.

$$H_k(\mathbb{K}, \mathbb{Z}_2) = \frac{\text{Ker } \partial_k}{\text{Im } \partial_{k+1}} = \frac{Z_k}{B_k}.$$

The inclusion map $f : \mathbb{K}_{i-1} \hookrightarrow \mathbb{K}_i$ defined by $f(x) = x$ induces a homomorphism $f_* : H_p(\mathbb{K}_{i-1}) \rightarrow H_p(\mathbb{K}_i)$. The nested sequence of complexes corresponds to a sequence of homology groups connected by the induced maps,

$$0 = H_p(\mathbb{K}_{-1}) \rightarrow H_p(\mathbb{K}_0) \rightarrow \cdots \rightarrow H_p(\mathbb{K}_n) = H_p(\mathbb{K})$$

Persistent homology studies how the homology groups change over the filtration.

Definition (positive simplex)

Given a filtration of \mathbb{K} , suppose $\mathbb{K}_i - \mathbb{K}_{i-1} = \sigma_i$, where σ_i is a $(k+1)$ -simplex. We call σ_i is **positive** if it belongs to a $(k+1)$ -cycle in \mathbb{K}_i and **negative** otherwise.

A positive simplex is also called a **generator**, a negative simplex a **killer**.

Definition (Betti Number)

Given a complex K , the i -th Betti number β_i is the rank of $H_i(K)$,

$$\beta_i = \text{Rank}H_i(K, \mathbb{Z}_2)$$

Suppose the number of positive k -simplexes is pos_k , and the number of negative k -simplexes is neg_k , then

$$\beta_k = \text{pos}_k - \text{neg}_{k+1}$$

Definition (Persistent Homology)

Define Z_k^l, B_k^l be the K -th cycle group and k -th boundary group respectively, of the l -complex K^l in a filtration. The p -persistent k -th homology group K^l is

$$H_k^{l,p} := \frac{Z_k^l}{B_k^{l+p} \cap Z_k^l}.$$

The p -persistent k -th Betti number $\beta_k^{l,p}$ of K^l is the rank of $H_k^{l,p}$.

Lemma

Consider the homomorphism $\eta_k^{l,p} : H_k^l \rightarrow H_k^{l+p}$, then

$$\text{img } \eta_k^{l,p} \cong H_k^{l,p}$$

Lemma

For each *positive* k -simplex σ^i , there exists a non-exact k -cycle c^i , c^i contains σ^i but no other positive k -simplices.

Proof.

Start with an arbitrary a k -cycle that contains σ^i and remove other positive k -simplices by adding their corresponding k -cycles. This method succeeds because each added cycle contains only one positive k -simplex by inductive assumption. \square

We use σ^i to represent c^i , and in turn the homologous class $[c^i] = c^i + B_k$.

$$\sigma^i \rightarrow c^i \rightarrow [c^i] = c^i + B_k. \quad \sigma^i \sim c^i$$

We add $[c^i]$ to the basis of $H_k(\mathbb{K}^i)$.




Figure: Generator, positive simplex.

For each **negative** $(k+1)$ -simplex σ^j , its boundary $d = \partial_{k+1}\sigma^j$ is a k -cycle, and can be represented as the linear combination of the basis of $H_k(\mathbb{K}_{j-1})$,

$$[d] = \sum_g [c^g], \{c^g\} \text{ basis } H_k(\mathbb{K}_{j-1}),$$

each $[c^g]$ is represented by a positive k -simplex σ^g , $g < j$, that is not yet paired. The collection of positive non-paired k -simplices is denoted as $\Gamma = \Gamma(d)$,

$$\Gamma(d) := \left\{ \sigma^g : [d] = \sum_g [c^g], \sigma^g \sim c^g \right\}$$

Suppose the youngest positive simplex in $\Gamma(\partial_{k+1}\sigma^j)$ is σ^i , then we form the pair (σ^i, σ^j) , and remove $[c^i]$ from $H_k(\mathbb{K}_j)$.

$[c^i]$ is created by σ^i and killed by σ^j , the persistence life of the k -cycle $[c^i]$ is $j - i - 1$.

Example Filtration




Figure: Generators and killers.

Filtration

$v_0, v_1, v_2, v_3, v_4, v_5, e_0, e_1, e_2, e_3, e_4, e_5, f_0, f_1, f_2$

Relabel them as

$\sigma^0, \sigma^1, \sigma^2, \sigma^3, \sigma^4, \sigma^5, \sigma^6, \sigma^7, \sigma^8, \sigma^9, \sigma^{10}, \sigma^{11}, \sigma^{12}, \sigma^{13}, \sigma^{14}, \sigma^{15}$

Example Generators




Figure: Generators.

Example Killers




Figure: Killers.

Example Pairing




Figure: Generators and killers.

$$\begin{aligned}\partial_2 f_2 &= e_2 + e_5 + e_6 + e_8 = (e_5 + 2e_4 + 2e_3) + (e_2 + 2e_1 + 2e_0) + e_6 + e_8 \\ &= (e_5 + e_4 + e_3) + (e_2 + e_1 + e_0) + \partial_2(f_0 + f_1)\end{aligned}$$

Key Lemma

Definition (Collision Free Cycle)

A collision free cycle is one where the youngest positive simplex has not been paired (killed).

Lemma (Collision)

Given a filtration, $\mathbb{K}_j - \mathbb{K}_{j-1} = \sigma^j$, σ^i is the youngest positive simplex in $\Gamma(\partial_{k+1}\sigma^j)$. Let e be a collision free k -cycle in \mathbb{K}_{j-1} homologous to $\partial_{k+1}\sigma^j$. Suppose the youngest positive simplex in e is σ^g , then

$$\sigma^i = \sigma^g.$$

$$\max \Gamma(\partial_{k+1}\sigma^j) = \max(e) \quad \forall e \text{ collision free, } [e] = [\partial\sigma^j].$$

Key Lemma

Proof.

Let f be the sum of the basis cycles, homologous to $d = \partial_{k+1}\sigma^j$. By definition, f 's youngest positive simplex is σ^i , namely the youngest simplex in $\Gamma(\partial_{k+1}\sigma^j)$,

$$\sigma^i = \max \Gamma(\partial_{k+1}\sigma^j).$$

This implies that there are no cycles homologous to d in \mathbb{K}_{i-1} or earlier complexes. Let σ^g be the youngest positive simplex in e . $[e] = [d]$, therefore $g \geq i$.

If $g > i$, then $e = f + c$, where c bounds in \mathbb{K}^{j-1} . $\sigma^g \notin f$, implies $\sigma^g \in c$, and as σ^g is the youngest in e , it is also the youngest in c . \square

continued.

Since e is collision free, the cycle created by σ^g , denoted as c^g , is still a non-boundary cycle in \mathbb{K}_{j-1} . Hence c^g can't be c , and can't be homologous to c when c becomes a boundary. Namely, when c is killed, σ^g is not paired yet.

It follows that the negative $(k+1)$ -simplex that kills c must pair a positive k -simplex in c , which is younger than σ^g , a contradiction.

This lemma shows, when σ^j is added to \mathbb{K}_{j-1} , we need to find any collision free cycle e homologous to $\partial_{k+1}\sigma^j$, and pair σ^j with the youngest positive simplex of e .

Pair Algorithm



Pair(σ)

- 1 $c = \partial_p \sigma$
- 2 τ is the youngest positive $(p - 1)$ -simplex in c .
- 3 **while** τ is paired and c is not empty **do**
- 4 find (τ, d) , d is the p -simplex paired with τ ;
- 5 $c \leftarrow \partial_p d + c$
- 6 Update τ to be the youngest positive $(p - 1)$ -simplex in c
- 7 **end while**
- 8 **if** c is not empty **then**
- 9 σ is negative p -simplex and paired with τ
- 10 **else**
- 11 σ is a positive p -simplex
- 12 **endif**

Handle Loop and Tunnel Loop



- 1 The simplices on the surface M are added into the filtration in any arbitrary order. Since $H_1(M)$ is of rank $2g$, the algorithm Pair generates $2g$ number of unpaired positive edges.
- 2 The simplices up to dimension 2 in I are added into the filtration. Since $H_1(I)$ of rank g , half of $2g$ positive edges generated in step 1 get paired with the negative triangles in I . Each pair corresponds to a killed loop, these g loops are handle loops.
- 3 Or the simplices up to dimension 2 in O are added into the filtration. Since $H_1(O)$ of rank g , half of $2g$ positive edges generated in step 2 get paired with the negative triangles in O . Each pair corresponds to a killed loop, these g loops are tunnel loops.

Example




$$3. \partial st = s + t, (t_1, st_3)$$


Example



Example



$$6. \partial sw = s + w, (w_5, sw_6)$$




$$7. tw$$


$$\begin{aligned} 7 : \partial tw7 &= w5 + t1 = w5 + t1 + \partial sw6 \\ &= w5 + t1 + (s0 + w5) = t1 + s0 + \partial st3 \\ &= t1 + s0 + (t1 + s0) \\ &= 0 \end{aligned}$$

$$c7 = (tw7, st3, sw6)$$

Example




$$8. \partial uv = u + v, (v_4, uv_8)$$




$$9. (u_2, sv_9)$$

$$\begin{aligned} 9. \partial sv_9 &= s_0 + v_4 = s_0 + v_4 + \partial uv_4 \\ &= s_0 + v_4 + (u_2 + v_4) \\ &= s_0 + u_2 \end{aligned}$$

Example



10. su_{10}




11. uw_{11}

Example


$$\begin{aligned}10. \partial su10 &= s0 + u2 \\ &= s0 + u2 + \partial sv9 \\ &= s0 + u2 + (s0 + v4) \\ &= u2 + v4 \\ &= u2 + v4 + \partial uv8 \\ &= u2 + v4 + (u2 + v4) \\ &= 0 \\ c10 &= (su10, uv8, sv9)\end{aligned}$$

$$\begin{aligned}11. \partial uw11 &= u2 + w5 \\ &= u2 + w5 + \partial sw6 \\ &= u2 + w5 + (s0 + w5) \\ &= s0 + u2 \\ &= s0 + u2 + \partial sv9 \\ &= s0 + u2 + (s0 + v4) \\ &= u2 + v4 \\ &= u2 + v4 + \partial uv8 \\ &= u2 + v4 + (u2 + v4) \\ &= 0. \\ c11 &= (uw11, uv8, sv9, sw6)\end{aligned}$$

Example



12. tu



13. $tuw, (tu_{12}, tuw_{13})$


Example

$$\begin{aligned}12. \partial tu_{12} &= t_1 + u_2 = t_1 + u_2 + \partial sv_9 \\ &= t_1 + u_2 + (s_0 + v_4) \\ &= t_1 + u_2 + s_0 + v_4 + \partial uv_8 \\ &= t_1 + u_2 + s_0 + v_4 + (u_2 + v_4) \\ &= t_1 + s_0 \\ &= s_0 + t_1 + \partial st_3 \\ &= s_0 + t_1 + (t_1 + s_0) \\ &= 0.\end{aligned}$$


$$c_{12} = (tu_{12}, st_3, sv_9, uv_8)$$

$$13. \partial tuw_{13} = tu_{12} + uw_{11} + wt_7 \\ (tuw_{13}, tu_{12})$$

Example



$$14. \partial suw = uw + su + sw \\ (uw_{11}, suw_{14})$$



$$15. stu, (tw_7, stu_{15})$$

Example

$$\begin{aligned} 15. \partial stu_{15} &= su_{10} + tu_{12} + st_3 \\ &= su_{10} + st_3 + tu_{12} + \partial tuw_{13} \\ &= su_{10} + st_3 + tu_{12} + (tu_{12} + uw_{11} + tw_7) \\ &= su_{10} + st_3 + uw_{11} + tw_7 \\ &= su_{10} + st_3 + uw_{11} + tw_7 + \partial suw_{14} \\ &= su_{10} + st_3 + uw_{11} + tw_7 + (sw_6 + su_{10} + uw_{11}) \\ &= st_3 + tw_7 + sw_6 \end{aligned}$$

Hence we obtain the pair (tw_7, stu_{15}) .

Example

$$\begin{aligned}17. \partial stw_{17} &= tw_7 + sw_6 + st_3 \\ &= sw_6 + st_3 + tw_7 + \partial stu_{15} \\ &= sw_6 + st_3 + tw_7 + (st_3 + tu_{12} + us_{10}) \\ &= sw_6 + tw_7 + (tu_{12} + us_{10}) + \partial tuw_{13} \\ &= sw_6 + tw_7 + (tu_{12} + us_{10}) + (tu_{12} + uw_{11} + wt_7) \\ &= sw_6 + us_{10} + uw_{11} \\ &= sw_6 + us_{10} + uw_{11} + \partial suw_{14} \\ &= sw_6 + us_{10} + uw_{11} + (su_{10} + uw_{11} + ws_6) \\ &= 0. \\ c_{17} &= (stw_{17}, stu_{15}, tuw_{13}, suw_{14}).\end{aligned}$$

Incidence Matrix

Assuming an ordering of the $(p - 1)$ simplices and of the p -simplices, the boundary of a p -chain can be obtained by multiplication of the corresponding vector with the incidence matrix,

$$\partial(c_p) = D_p c_p.$$

The incidence matrix is defined as

$$D_p[i, j] = \begin{cases} 1 & \sigma_i^{p-1} \in \sigma_j^p \\ 0 & \sigma_i^{p-1} \notin \sigma_j^p \end{cases}$$

Incidence Matrix and Betti Number

A classic algorithm computes the Betti numbers of K by reducing its incidence matrices to Smith normal form. It uses row and column operations to zero out all entries except along an initial portion of the diagonal.




Figure: Smith norm of incidence matrix in \mathbb{Z}_2 .

The Betti number

$$\beta_p = \text{rank} Z_p - \text{rank} B_p.$$

Pairing Algorithm

Definition (Monotonous Filtering)

A filtering is monotonous, if in the ordering of K , any simplex σ is preceded by its faces.

An algorithm computes the persistence diagrams by pairing the simplices, which uses column operator to reduce D and another $0 - 1$ matrix R . Let $\text{low}_R(j)$ be the row index of the last 1 in column j of R , and (undefined if the column is zero).

Definition (Reduced Matrix and Pairing)

We call R reduced and low_R a pairing function, if

$$\text{low}_R(j) \neq \text{low}_R(j'),$$

whenever $j \neq j'$ specify two non-zero columns.

Pairing Algorithm

Algorithm: Incidence matrix reduction

- 1 $R \leftarrow D$
- 2 **for** $j = 1$ **to** n **do**
- 3 **while** $\exists j' < j$ **with** $\text{low}_R(j') = \text{low}_R(j)$ **do**
- 4 add column j' to column j
- 5 **endwhile**
- 6 **endfor.**

The pairing is given by

$$(\sigma_i, \sigma_j) \iff i = \text{low}_R(j).$$

σ_i is positive, it generates a homology class; σ_j is negative, it kills a homology class.




Figure: Generators and killers.

Pairing by matrix induction

Boundary operator ∂_1 , incidence matrix D_1 ,

	e_0	e_1	e_2	e_3	e_4	e_5	e_6	e_7	e_8
v_0	1	0	1	0	0	0	1	0	0
v_1	1	1	0	0	0	0	0	1	0
v_2	0	1	1	0	0	0	0	0	1
v_3	0	0	0	1	0	1	1	0	0
v_4	0	0	0	1	1	0	0	1	0
v_5	0	0	0	0	1	1	0	0	1

Pairing by matrix induction

$1 + 2, 4 + 5, 3 + 7, 4 + 8$

	e_0	e_1	e_2	e_3	e_4	e_5	e_6	e_7	e_8
v_0	1	0	1	0	0	0	1	0	0
v_1	1	1	1	0	0	0	0	1	0
v_2	0	1	0	0	0	0	0	0	1
v_3	0	0	0	1	0	1	1	1	0
v_4	0	0	0	1	1	1	0	0	1
v_5	0	0	0	0	1	0	0	0	0

Pairing by matrix induction

$1 + 2, 3 + 5, 3 + 8$

	e_0	e_1	e_2	e_3	e_4	e_5	e_6	e_7	e_8
v_0	1	0	0	0	0	0	1	0	0
v_1	1	1	0	0	0	0	0	1	0
v_2	0	1	0	0	0	0	0	0	1
v_3	0	0	0	1	0	0	1	1	1
v_4	0	0	0	1	1	0	0	0	0
v_5	0	0	0	0	1	0	0	0	0

Pairing by matrix induction

$6 + 7, 6 + 8$

	e_0	e_1	e_2	e_3	e_4	e_5	e_6	e_7	e_8
v_0	1	0	0	0	0	0	1	1	1
v_1	1	1	0	0	0	0	0	1	0
v_2	0	1	0	0	0	0	0	0	1
v_3	0	0	0	1	0	0	1	0	0
v_4	0	0	0	1	1	0	0	0	0
v_5	0	0	0	0	1	0	0	0	0

Pairing by matrix induction

$$0 + 7, 1 + 8, 0 + 8$$

	e_0	e_1	e_2	e_3	e_4	e_5	e_6	e_7	e_8
v_0	1	0	0	0	0	0	1	0	0
v_1	1	1	0	0	0	0	0	0	0
v_2	0	1	0	0	0	0	0	0	0
v_3	0	0	0	1	0	0	1	0	0
v_4	0	0	0	1	1	0	0	0	0
v_5	0	0	0	0	1	0	0	0	0

Generators e_2, e_5, e_7, e_8 , corresponding to 0 columns. Killers corresponds to non-zero columns. Pairing

$$(e_0, v_1), (e_1, v_2), (e_3, v_4), (e_4, v_5), (e_6, v_3)$$

Pairing by matrix induction

	f_0	f_1	f_2
e_0	1	0	0
e_1	0	1	0
e_2	0	0	1
e_3	1	0	0
e_4	0	1	0
e_5	0	0	1
e_6	1	0	1
e_7	1	1	0
e_8	0	1	1

$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix} \xrightarrow{2+3} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 1 \\ 0 & 1 & 0 \end{bmatrix} \xrightarrow{1+3} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

The pairing is

$$(f_0, e_7), (f_1, e_8), (f_2, e_5)$$

Topological Annulus






Figure: Topological Annulus.

Topological Torus



Topological Torus



Topological Torus




Figure: Handle and tunnel loops.

Topological Torus




Figure: Handle and tunnel loops.

Topological Torus




Figure: Interior and exterior volumes.

Topological Quadrilaterals






Figure: Quadrilateral example.

Topological Quadrilaterals




$$\partial f_5 = c_{17} = e_{15} + e_9 + e_8 + e_{17}, \quad \partial f_1 = c_{19} = e_3 + e_2 + e_{14} + e_{19}.$$

Topological Quadrilaterals




$$\partial f_3 = c_{18} = e_{16} + e_5 + e_6 + e_{18}, \quad \partial f_0 = c_{23} = e_{12} + e_{14} + e_1 + e_{23}.$$

Topological Quadrilaterals




$$c_{21} = e_{16} + e_4 + e_3 + e_2 + e_{14} + e_{21}, \quad \partial f_2 = c_{19} + c_{21}$$

Topological Quadrilaterals



$$c_{22} = e_{15} + e_4 + e_{16} + e_9 + e_5 + e_8 + e_7 + e_6 + e_{22}, \quad \partial f_4 = c_{22} + c_{18} + c_{17}$$


Topological Quadrilaterals



$$c_{13} = e_{12} + e_{10} + e_9 + e_8 + e_7 + e_6 + e_5 + e_4 + e_3 + e_2 + e_1 + e_{13}, \partial f_7 = c_{11} + c_{13}$$


$$c_{11} = e_0 + e_1 + e_2 + e_3 + e_4 + e_5 + e_6 + e_7 + e_8 + e_9 + e_{10} + e_{11}$$

Topological Quadrilaterals




$$c_{20} = e_{15} + e_9 + e_8 + e_7 + e_6 + e_5 + e_4 + e_3 + e_2 + e_2 + e_{12} + e_{20}, \partial f_6 = c_{20} + c_{13}.$$

Topological Quadrilaterals



$$\begin{aligned} \partial f_0 &= c_{23}, (e_{23}, f_0), \\ \partial f_1 &= c_{19}, (e_{19}, f_1), \\ \partial f_2 &= c_{21} + \partial f_1, (e_{21}, f_2), \\ \partial f_3 &= c_{18}, (e_{18}, f_3), \\ \partial f_5 &= c_{17}, (e_{17}, f_5), \\ \partial f_4 &= \\ & c_{17} + c_{18} + c_{22}, (e_{22}, f_4), \\ \partial f_6 &= c_{13} + c_{20}, (e_{20}, f_6), \\ \partial f_7 &= c_{13} + c_{11}, (e_{13}, f_7), \end{aligned}$$

Topological Quadrilaterals



$$\partial f_8 \rightarrow \partial f_0, \partial f_4, \partial f_2, \partial f_6, \partial f_1, \partial f_3, \partial f_5, \partial f_7$$

$$\rightarrow e_0 + e_1 + e_2 + e_3 + e_4 + e_5 + e_6 + e_7 + e_8 + e_9 + e_{10} + e_{11}$$