

# de Rham Cohomology, Hodge Decomposition

David Gu

Yau Mathematics Science Center  
Tsinghua University  
Computer Science Department  
Stony Brook University

*gu@cs.stonybrook.edu*

July 13, 2022

# Exterior Differential

The homology of a manifold is the difference between the **closed loops** and the **boundary loops**.

The cohomology of a manifold is the difference between the **curl free** vector fields and the **gradient** vector fields.

Consider a planar vector field defined on  $\mathbb{C} \setminus \{0\}$ ,

$$\mathbf{v}(x, y) = \left( \frac{-y}{x^2 + y^2}, \frac{x}{x^2 + y^2} \right)$$

direct computation  $\nabla \times \mathbf{v}(x, y) = 0$ .

$$\left| \begin{array}{cc} \frac{\partial}{\partial x} & \frac{\partial}{\partial y} \\ \frac{-y}{x^2+y^2} & \frac{x}{x^2+y^2} \end{array} \right| = 0.$$

But choose the unit circle

$$\oint_{\gamma} \omega = \oint_{\gamma} d \tan^{-1} \frac{y}{x} = 2\pi$$

therefore  $\mathbf{v}$  is not a gradient field. Namely,  $d\theta$  locally is integrable, globally not.

# Smooth Manifold




Figure: A manifold.

## Definition (Manifold)

A manifold is a topological space  $M$  covered by a set of open sets  $\{U_\alpha\}$ . A homeomorphism  $\phi_\alpha : U_\alpha \rightarrow \mathbb{R}^n$  maps  $U_\alpha$  to the Euclidean space  $\mathbb{R}^n$ .  $(U_\alpha, \phi_\alpha)$  is called a coordinate chart of  $M$ . The set of all charts  $\{(U_\alpha, \phi_\alpha)\}$  form the atlas of  $M$ . Suppose  $U_\alpha \cap U_\beta \neq \emptyset$ , then

$$\phi_{\alpha\beta} = \phi_\beta \circ \phi_\alpha^{-1} : \phi_\alpha(U_\alpha \cap U_\beta) \rightarrow \phi_\beta(U_\alpha \cap U_\beta)$$

is a transition map.

If all transition maps  $\phi_{\alpha\beta} \in C^\infty(\mathbb{R}^n)$  are smooth, then the manifold is a differential manifold or a smooth manifold.

# Tangent Space

## Definition (Tangent Vector)

A tangent vector  $\xi$  at the point  $p$  is an association to every coordinate chart  $(x^1, x^2, \dots, x^n)$  at  $p$  an  $n$ -tuple  $(\xi^1, \xi^2, \dots, \xi^n)$  of real numbers, such that if  $(\tilde{\xi}^1, \tilde{\xi}^2, \dots, \tilde{\xi}^n)$  is associated with another coordinate system  $(\tilde{x}^1, \tilde{x}^2, \dots, \tilde{x}^n)$ , then it satisfies the transition rule

$$\tilde{\xi}^i = \sum_{j=1}^n \frac{\partial \tilde{x}^i}{\partial x^j}(p) \xi^j.$$

A smooth vector field  $\xi$  assigns a tangent vector for each point of  $M$ , it has local representation

$$\xi(x^1, x^2, \dots, x^n) = \sum_{i=1}^n \xi_i(x^1, x^2, \dots, x^n) \frac{\partial}{\partial x_i}.$$

$\left\{ \frac{\partial}{\partial x_i} \right\}$  represents the vector fields of the velocities of iso-parametric curves on  $M$ . They form a basis of all vector fields.

## Definition (Push-forward)

Suppose  $\phi : M \rightarrow N$  is a differential map from  $M$  to  $N$ ,  $\gamma : (-\epsilon, \epsilon) \rightarrow M$  is a curve,  $\gamma(0) = p$ ,  $\gamma'(0) = \mathbf{v} \in T_p M$ , then  $\phi \circ \gamma$  is a curve on  $N$ ,  $\phi \circ \gamma(0) = \phi(p)$ , we define the tangent vector

$$\phi_*(\mathbf{v}) = (\phi \circ \gamma)'(0) \in T_{\phi(p)} N,$$

as the push-forward tangent vector of  $\mathbf{v}$  induced by  $\phi$ .



## Definition (Differential 1-form)

The tangent space  $T_p M$  is an  $n$ -dimensional vector space, its dual space  $T_p^* M$  is called the cotangent space of  $M$  at  $p$ . Suppose  $\omega \in T_p^* M$ , then  $\omega : T_p M \rightarrow \mathbb{R}$  is a linear function defined on  $T_p M$ ,  $\omega$  is called a differential 1-form at  $p$ .

A differential 1-form field has the local representation

$$\omega(x^1, x^2, \dots, x^n) = \sum_{i=1}^n \omega_i(x^1, x^2, \dots, x^n) dx_i,$$

where  $\{dx_i\}$  are the differential forms dual to  $\{\frac{\partial}{\partial x_j}\}$ , such that

$$\langle dx_i, \frac{\partial}{\partial x_j} \rangle = dx_i \left( \frac{\partial}{\partial x_j} \right) = \delta_{ij}.$$

# High order exterior forms

## Definition (Tensor)

A tensor  $\Theta$  of type  $(m, n)$  on a manifold  $M$  is a correspondence that associates to each point  $p \in M$  a multi-linear map

$$\Theta_p : T_p M \times T_p M \times \cdots \times T_p^* M \cdots \times T_p^* M \rightarrow \mathbb{R},$$

where the tangent space  $T_p M$  appears  $m$  times and cotangent space  $T_p^* M$  appears  $n$  times.

## Definition (exterior $m$ -form)

An exterior  $m$ -form is a tensor  $\omega$  of type  $(m, 0)$ , which is skew symmetric in its arguments, namely

$$\omega_p(\xi_{\sigma(1)}, \xi_{\sigma(2)}, \cdots, \xi_{\sigma(m)}) = (-1)^\sigma \omega_p(\xi_1, \xi_2, \cdots, \xi_m)$$

for any tangent vectors  $\xi_1, \xi_2, \cdots, \xi_m \in T_p M$  and any permutation  $\sigma \in S_m$ , where  $S_m$  is the permutation group.

## Differential Form

The local representation of  $\omega$  in  $(x^1, x^2, \dots, x^m)$  is

$$\omega = \sum_{1 \leq i_1 < i_2 < \dots < i_m \leq n} \omega_{i_1 i_2 \dots i_m} dx^{i_1} \wedge dx^{i_2} \wedge \dots \wedge dx^{i_m} = \omega_I dx^I,$$

$\omega_I$  is a function of the reference point  $p$ ,  $\omega$  is said to be differentiable, if each  $\omega_I$  is differentiable.

## Definition (Wedge product)

A coordinate free representation of wedge product of  $m_1$ -form  $\omega_1$  and  $m_2$ -form  $\omega_2$  is defined as  $(\omega_1 \wedge \omega_2)(\xi_1, \xi_2, \dots, \xi_{m_1+m_2})$  equals

$$\sum_{\sigma \in S_{m_1+m_2}} \frac{(-1)^\sigma}{m_1! m_2!} \omega_1(\xi_{\sigma(1)}, \dots, \xi_{\sigma(m_1)}) \omega_2(\xi_{\sigma(m_1+1)}, \dots, \xi_{\sigma(m_1+m_2)})$$

# Wedge product

Give  $k$  differential 1-forms, their exterior wedge product is given by:

$$\omega_1 \wedge \omega_2 \cdots \omega_k(v_1, v_2, \dots, v_k) = \begin{vmatrix} \omega_1(v_1) & \omega_1(v_2) & \cdots & \omega_1(v_k) \\ \omega_2(v_1) & \omega_2(v_2) & \cdots & \omega_2(v_k) \\ \vdots & \vdots & \ddots & \vdots \\ \omega_k(v_1) & \omega_k(v_2) & \cdots & \omega_k(v_k) \end{vmatrix}$$

Exterior is anti-symmetric, suppose  $\sigma \in S_k$  is a permutation, then

$$\omega_{\sigma(1)} \wedge \omega_{\sigma(2)} \wedge \cdots \wedge \omega_{\sigma(k)} = (-1)^\sigma \omega_1 \wedge \omega_2 \wedge \cdots \wedge \omega_k.$$

## Definition (Pull back)

Suppose  $\phi : M \rightarrow N$  is a differentiable map from  $M$  to  $N$ ,  $\omega$  is an  $m$ -form on  $N$ , then the pull-back  $\phi^*\omega$  is an  $m$ -form on  $M$  defined by

$$(\phi^*\omega)_p(\xi_1, \dots, \xi_m) = \omega_{\phi(p)}(\phi_*\xi_1, \dots, \phi_*\xi_m), p \in M$$

for  $\xi_1, \xi_2, \dots, \xi_m \in T_pM$ , where  $\phi_*\xi_j \in T_{\phi(p)}N$  is the push forward of  $\xi_j \in T_pM$ .


## Integration in Euclidean space

Suppose that  $U \subset \mathbb{R}^n$  is an open set,

$$\omega = f(x)dx^1 \wedge dx^2 \wedge \cdots \wedge dx^n,$$

then

$$\int_U \omega = \int_U f(x)dx^1 dx^2 \cdots dx^n.$$




Suppose  $U \subset M$  is an open set of a manifold  $M$ , a chart  $\phi : U \rightarrow \Omega \subset \mathbb{R}^n$ , then

$$\int_U \omega = \int_{\Omega} (\phi^{-1})^* \omega.$$



# Integration



Integration is independent of the choice of the charts. Let  $\psi : U \rightarrow \psi(U)$  be another chart, with local coordinates  $(u_1, u_2, \dots, u_n)$ , then

$$\int_{\phi(U)} f(x) dx^1 dx^2 \dots dx^n = \int_{\psi(U)} f(x(u)) \det \left( \frac{\partial x^i}{\partial u^j} \right) du^1 du^2 \dots du^n.$$

## Integration on Manifolds

consider a covering of  $M$  by coordinate charts  $\{(U_\alpha, \phi_\alpha)\}$  and choose a partition of unity  $\{f_i\}$ ,  $i \in I$ , such that  $f_i(p) \geq 0$ ,

$$\sum_i f_i(p) \equiv 1, \forall p \in M.$$

Then  $\omega_i = f_i\omega$  is an  $n$ -form on  $M$  with compact support in some  $U_\alpha$ , we can set the integration as

$$\int_M \omega = \sum_i \int_M \omega_i.$$

# Exterior Derivative

## Exterior Derivative of a Function

Suppose  $f : M \rightarrow \mathbb{R}$  is a differentiable function, then the exterior derivative of  $f$  is a 1-form,

$$df = \sum_i \frac{\partial f}{\partial x_i} dx^i.$$

## Exterior Derivative of Differential Forms

The exterior derivative of an  $m$ -form on  $M$  is an  $(m + 1)$ -form on  $M$  defined in local coordinates by

$$d\omega_I = d(\omega_I dx^I) = (d\omega_I) \wedge dx^I,$$

where  $d\omega_I$  is the differential of the function  $\omega_I$ .

The exterior derivative of a differential 1-form is given by:

$$d\left(\sum \omega_i dx_i\right) = \sum_{i,j} \left(\frac{\partial \omega_j}{\partial x_i} - \frac{\partial \omega_i}{\partial x_j}\right) dx_i \wedge dx_j,$$

that of a differential  $k$ -form

$$d(\omega_1 \wedge \omega_2 \cdots \wedge \omega_k) = \sum (-1)^{i-1} \omega_1 \wedge \cdots \wedge \omega_{i-1} \wedge d\omega_i \wedge \omega_{i+1} \wedge \cdots \wedge \omega_k.$$

## Theorem (Stokes)

let  $M$  be an  $n$ -manifold with boundary  $\partial M$  and  $\omega$  be a differentiable  $(n - 1)$ -form with compact support on  $M$ , then

$$\int_{\partial M} \omega = \int_M d\omega.$$

# Stokes Theorem

## Theorem

Suppose  $\Sigma$  is a differential manifold, then we have

$$d^k \circ d^{k-1} = 0.$$

## Proof.

Assume  $\omega$  is a  $k - 1$  differential form,  $D$  is a  $k + 1$  chain, from Stokes theorem, we have

$$\int_D d^k \circ d^{k-1} \omega = \int_{\partial_{k+1} D} d^{k-1} \omega = \int_{\partial_k \circ \partial_{k+1} D} \omega = 0,$$

since  $\partial_k \circ \partial_{k+1} = 0$ . □

Let  $\Omega^k(\Sigma)$  be the space of all differential  $k$ -forms,  $d^k : \Omega^k(\Sigma) \rightarrow \Omega^{k+1}(\Sigma)$  be exterior differential operator.

## Definition (Closed form)

$k$ -form  $\omega \in \Omega^k(\Sigma)$  is called a closed form, if  $d^k\omega = 0$ , namely  $\omega \in \text{Ker } d^k$ .

## Definition (Exact Form)

$k$ -differential form  $\omega \in \Omega^k(\Sigma)$  is called exact form, if there is a  $\tau \in \Omega^{k-1}(\Sigma)$ , such that  $\omega = d^{k-1}\tau$ , namely  $\omega \in \text{Im } d^{k-1}$ .

Since  $d^k \circ d^{k-1} = 0$ , exact forms are closed,  $\text{Im } d^{k-1} \subset \text{Ker } d^k$ .

## Definition (de Rham Cohomology)

Assume  $\Sigma$  is a differential manifold, then de Rham complex is

$$\Omega^0(\Sigma, \mathbb{R}) \xrightarrow{d^0} \Omega^1(\Sigma, \mathbb{R}) \xrightarrow{d^1} \Omega^2(\Sigma, \mathbb{R}) \xrightarrow{d^2} \Omega^3(\Sigma, \mathbb{R}) \xrightarrow{d^3} \dots$$

$$H_{dR}^k(\Sigma, \mathbb{R}) := \frac{\text{Ker } d^k}{\text{Im } d^{k-1}}$$

## Theorem

*The de Rham cohomology group  $H_{dR}^m(M)$  is isomorphic to the cohomology group  $H^m(M, \mathbb{R})$*

$$H_{dR}^m(M) \cong H^m(M, \mathbb{R}).$$



# Hodge Operator




Figure: Hodge star operator.

# Hodge Star Operator - First Definition


Suppose  $M$  is a Riemannian manifold, we can locally find oriented orthonormal basis of vector fields, and choose parameterization, such that

$$\left\{ \frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \dots, \frac{\partial}{\partial x_n} \right\}$$

form an oriented orthonormal basis. let

$$\{dx_1, dx_2, \dots, dx_n\}$$

be the dual 1-form basis.



## Definition (Hodge Star Operator)

The Hodge star operator  $*$  :  $\Omega^k(M) \rightarrow \Omega^{n-k}(M)$  is a linear operator

$$*(dx_1 \wedge dx_2 \wedge \cdots \wedge dx_k) = dx_{k+1} \wedge dx_{k+2} \wedge \cdots \wedge dx_n.$$

## Hodge Star Operator

Let  $\sigma = (i_1, i_2, \dots, i_n)$  be a permutation, then the Hodge star operator

$$*(dx_{i_1} \wedge dx_{i_2} \wedge \cdots \wedge dx_{i_k}) = (-1)^\sigma dx_{i_{k+1}} \wedge dx_{i_{k+2}} \wedge \cdots \wedge dx_{i_n}.$$

## Definition

Let  $\eta, \zeta \in \Omega^k(M)$  are two  $k$ -forms on  $M$ , then the norm is defined as

$$(\eta, \zeta) = \int_M \eta \wedge * \zeta.$$

$\Omega^k(M)$  is a Hilbert space.

# Hodge Star Operator - Second Equivalent Definition

Given a Riemannian manifold  $(M, \mathbf{g})$ ,  $\mathbf{g} = (g_{ij})$ , which gives the inner product in the tangent space  $T_p(M)$ ,

$$g_{ij} = \langle \partial_i, \partial_j \rangle_{\mathbf{g}}.$$

its inverse matrix is  $(g^{ij})$ , satisfies

$$\sum_{j=1}^n g_{ij} g^{jk} = \delta_i^k.$$

## Definition (Dual Inner Product)

Given a  $n$  dimensional Riemannian manifold  $(M, \mathbf{g})$ , the dual inner product  $\langle \cdot, \cdot \rangle_{\mathbf{g}} : T_p^*(M) \times T_p^*(M) \rightarrow \mathbb{R}$ ,  $\forall \omega, \eta \in T_p^*(M)$ ,  $\omega = \sum_{i=1}^n \omega_i dx^i$ ,  $\eta = \sum_{i=1}^n \eta_i dx^i$ , then

$$\langle \omega, \eta \rangle_{\mathbf{g}} = \sum_{i,j=1}^n g^{ij} \omega_i \eta_j.$$

## Orthonormal Basis

Let  $\{\theta_1, \theta_2, \dots, \theta_n\}$  is a set of orthonormal basis

$$\langle \theta_i, \theta_j \rangle_{\mathbf{g}} = \delta_i^j.$$

## Basis of $\Omega^k(M)$

We use  $\{\theta_i\}$  to construct the basis of  $\Omega^k(M)$ ,

$$\Omega^k(M) := \text{Span}\{\theta_{i_1} \wedge \theta_{i_2} \wedge \dots \wedge \theta_{i_k} \mid i_1 < i_2 < \dots < i_k\}.$$

## Dual Inner Product

We define dual inner product  $\langle \cdot, \cdot \rangle_{\mathbf{g}} : \Omega^k(M) \times \Omega^k(M)$  as follows:

$$\langle \theta_{i_1} \wedge \dots \wedge \theta_{i_k}, \theta_{j_1} \wedge \dots \wedge \theta_{j_k} \rangle = \delta_{i_1 \dots i_k}^{j_1 \dots j_k}.$$



# Riemannian Volume Element

## Riemannian volume Element

Let  $G = \det(g_{ij})$ , then in the local coordinates, the Riemannian volume element is defined as

$$\omega_{\mathbf{g}} = \sqrt{G} dx^1 \wedge dx^2 \wedge \cdots \wedge dx^n.$$

## Definition (Hodge Star Operator)

$$* : \Omega^k(M) \rightarrow \Omega^{n-k}(M),$$

$$\omega \wedge *\eta = \langle \omega, \tau \rangle_{\mathbf{g}} \omega_{\mathbf{g}}.$$

Therefore

$$*(1) = \omega_{\mathbf{g}}, \quad *\omega_{\mathbf{g}} = 1.$$

## Definition (Inner Product)

Let  $(M, \mathbf{g})$  be a  $n$  dimensional Riemannian manifold,  $\zeta$  and  $\eta$  are differential  $k$ -forms,  $0 \leq k \leq n$ , then  $\zeta$  and  $\eta$  inner product is defined as

$$(\zeta, \eta) := \int_M \zeta \wedge \ast \eta = \int_M \langle \zeta, \eta \rangle_{\mathbf{g}} \omega_{\mathbf{g}}$$

# Hodge Star Operator on Surface - Type I

Suppose  $(S, \mathbf{g})$  is a surface with a Riemannian metric, with isothermal coordinates  $(u, v)$ , the metric is

$$\mathbf{g} = e^{2\lambda(u,v)}(du^2 + dv^2),$$

Then

$$\frac{\partial}{\partial x_1} = e^{-\lambda} \frac{\partial}{\partial u}, \quad \frac{\partial}{\partial x_2} = e^{-\lambda} \frac{\partial}{\partial v},$$

And

$$dx_1 = e^\lambda du, \quad dx_2 = e^\lambda dv.$$

Hodge Star

$$*dx_1 = dx_2, \quad *du = dv$$

$$*dx_2 = -dx_1, \quad *dv = -du$$

$$*(1) = dx_1 \wedge dx_2 = e^2 du \wedge dv, \quad *(dx_1 \wedge dx_2) = 1.$$

# Hodge Star Operator on Surface - Type II

Suppose  $(S, \mathbf{g})$  is a surface with a Riemannian metric, with isothermal coordinates  $(u, v)$ , the metric is

$$\mathbf{g} = e^{2\lambda(u,v)}(du^2 + dv^2),$$

surface area element is

$$\omega_{\mathbf{g}} = e^{2\lambda(u,v)} du \wedge dv.$$

Given 1-forms  $\omega = \omega_1 du + \omega_2 dv$  and  $\tau = \tau_1 du + \tau_2 dv$ , its wedge product is

$$\omega \wedge \tau = (\omega_1 \tau_2 - \omega_2 \tau_1) du \wedge dv.$$

Inner product is

$$\langle \omega, \tau \rangle_{\mathbf{g}} = e^{-2\lambda(u,v)} (\omega_1 \tau_1 + \omega_2 \tau_2).$$

# Hodge Star Operator on Surface

$$(\omega_1 du + \omega_2 dv) \wedge *du = \langle \omega, du \rangle_{\mathbf{g}} \omega_{\mathbf{g}} = e^{-2\lambda} \omega_1 e^{2\lambda} du \wedge dv,$$

This shows  $*du = dv$ , similarly  $*dv = -du$ .

$$*(\omega_1 du + \omega_2 dv) = \omega_1 dv - \omega_2 du.$$

Hence  $**\omega = -\omega$ .

# Hodge Decomposition

# Codifferential operator

## Definition

The codifferential operator  $\delta : \Omega^k(M) \rightarrow \Omega^{k-1}(M)$  is defined as

$$\delta = (-1)^{kn+n+1} d^*,$$

where  $d$  is the exterior derivative.

## Lemma

*The codifferential is the adjoint of the exterior derivative, in that*

$$(\delta\zeta, \eta) = (\zeta, d\eta).$$

# Laplace Operator

## Definition (Laplace Operator)

The Laplace operator  $\Delta : \Omega^k(M) \rightarrow \Omega^k(M)$ ,

$$\Delta = d\delta + \delta d.$$

## Lemma

*The Laplace operator is symmetric*

$$(\Delta\zeta, \eta) = (\zeta, \Delta\eta)$$

*and non-negative*

$$(\Delta\eta, \eta) \geq 0.$$

## Proof.

$$(\Delta\zeta, \eta) = (d\zeta, d\eta) + (\delta\zeta, \delta\eta).$$



# Harmonic Forms

## Definition (Harmonic forms)

Suppose  $\omega \in \Omega^k(M)$ , then  $\omega$  is called a  $k$ -harmonic form, if

$$\Delta\omega = 0.$$

## Lemma

$\omega$  is a harmonic form, if and only if

$$d\omega = 0, \delta\omega = 0.$$

## Proof.

$$0 = (\Delta\omega, \omega) = (d\omega, d\omega) + (\delta\omega, \delta\omega).$$




# Hodge Decomposition

## Definition (Harmonic form group)

All harmonic  $k$ -forms form a group, denoted as  $H_{\Delta}^k(M)$ .

## Theorem (Hodge Decomposition)

$$\Omega_k = \text{img} d^{k-1} \oplus \text{img} \delta^{k+1} \oplus H_{\Delta}^k(M).$$



# Hodge Decomposition

Proof.

$(\text{img}d^{k-1})^\perp = \{\omega \in \Omega^k(M) \mid (\omega, d\eta) = 0, \forall \eta \in \Omega^{k-1}(M)\}$ , because  $(\omega, d\eta) = (\delta\omega, \eta)$ , so  $(\text{img}d^{k-1})^\perp = \ker\delta^k$ . similarly,  $(\text{img}\delta^{k+1})^\perp = \ker d^k$ . Because  $\text{img}d^{k-1} \subset \ker d^k$ ,  $\text{img}\delta^{k+1} \subset \ker\delta^k$ , therefore  $\text{img}d^{k-1} \perp \text{img}\delta^{k+1}$ ,

$$\Omega^k = \text{img}d^{k-1} \oplus \text{img}\delta^{k+1} \oplus (\text{img}d^{k-1} \oplus \text{img}\delta^{k+1})^\perp$$

$$(\text{img}d^{k-1} \oplus \text{img}\delta^{k+1})^\perp = (\text{img}d^{k-1})^\perp \cap (\text{img}\delta^{k+1})^\perp = \ker\delta^k \cap \ker d^k = H_\Delta^k. \quad \square$$

# Hodge Decomposition

suppose  $\omega \in \ker d^k$ , then  $\omega \perp \operatorname{img} \delta^{k+1}$ , then  $\omega = \alpha + \beta$ ,  $\alpha \in \operatorname{img} d^{k-1}$ ,  $\beta \in H_{\Delta}^k(M)$ , define project  $h : \ker d^k \rightarrow H_{\Delta}^k(M)$ ,

## Theorem

Suppose  $\omega$  is a closed form, its harmonic component is  $h(\omega)$ , then the map:

$$h : H_{dR}^k(M) \rightarrow H_{\Delta}^k(M).$$

is isomorphic.

Each cohomologous class has a unique harmonic form.

# Harmonic 1-forms






Figure: Harmonic 1-form group basis.

# Dual Mesh



Delaunay triangulation  $T$  and Voronoi diagram  $D$ , every prime edge  $e$  corresponds to an dual edge  $\bar{e}$ .  $\omega$  is a one-form on  $T$ .  $*\omega$  is a one-form on  $D$ .


# Discrete Hodge Operator



Discrete Hodge star operator,

$$\frac{\omega(e)}{e} = \frac{* \omega(\bar{e})}{|\bar{e}|}, \quad * \omega(\bar{e}) = \frac{|\bar{e}|}{|e|} \omega(e) = \frac{1}{2}(\cot \alpha + \cot \beta) \omega(e).$$

# Discrete Harmonic One-form




$\omega$  is closed,

$$d\omega = 0.$$



# Discrete Harmonic One-form



$\omega$  is coclosed,

$$\delta\omega = *d^*\omega = 0.$$