

Adaptive Color Structured Light for Calibration and Shape Reconstruction

- Supplementary Materials -

Xin Dong^{*}
Southwest University

Haibin Ling[†]
Stony Brook University

Bingyao Huang[‡]
Southwest University

1 INTRODUCTION

In the supplementary materials, We provide more detailed experimental results and data. First, to analyze the methods of adaptive color SL and MAP more deeply, we conduct ablation studies and we will show more detailed experimental results in § 2. Then, to further verify the accuracy of the calibration, the reconstruction results are shown in § 3. Finally, we show the quantified results of the single-shot reconstruction in § 4. In addition to the presentation of the experimental results, we also briefly display different poses of the calibration board under the imaging setting and ambient light condition in Fig. 1 and Fig. 2.

2 ABLATION STUDIES

To verify the effectiveness of adaptive color SL and MAP more deeply, we come to the ablation experiment. In the ablation experiment, we conduct an experiment using Otsu [4] to perform color detection on the colors of the adaptive color SL, termed as **Ours w/o MAP**. Due to $\mathbf{Q}_{1:N}$ are obtained based on the generation process of adaptive color SL and MAP needs to use $\mathbf{Q}_{1:N}$, the fixed color SL does not have corresponding $\mathbf{Q}_{1:N}$ and MAP cannot be used to perform color detection on the colors of fixed color SL.

The comparisons of color detection and grid segmentation performance under different imaging settings and ambient light conditions for Huang [2], Ours w/o MAP and Ours are shown in Tab. 1 and Tab. 2.

Calibration RMS reprojection errors (pixels) of the camera, the projector and stereo under different imaging settings and ambient light conditions for Moreno & Taubin [3], Huang [2], Ours w/o MAP and Ours are shown in Tab. 3 and Tab. 4.

It should be noted that the performance of Ours w/o MAP varies greatly under different imaging settings and ambient light conditions. Otsu [4] relies on the characteristics of the color hue distribution, that is, when the intra-class variances of the hue distributions of different colors are very different, this can easily cause Otsu [4] to make mistakes, and please refer to [5] for detailed proof. The MAP no longer constrains the characteristics of the color hue distribution, as long as the color distinction of different colors is large.

3 RECONSTRUCTION

To further verify the accuracy of the calibration, the obtained calibration parameters of the four methods are also used to reconstruct point clouds for different real objects using Moreno & Taubin's [3] Gray-coded SL patterns.

The point cloud alignment errors (mm) of David (mm), girl (mm) and box (mm) for Moreno & Taubin [3], Huang [2], Ours w/o MAP and Ours under different imaging settings and ambient light conditions are shown in Tab. 5, Tab. 6, Tab. 7, Tab. 8, Tab. 9 and Tab. 10.




Figure 1: The different poses of the calibration board in Setting2.




Figure 2: The different poses of the calibration board in Light2.

4 SINGLE-SHOT RECONSTRUCTION

The methods that are applied to scan the object from different views (about 7-12 shots) and merge the point clouds using ICP [1] are suffixed with (**merged**). The number of reconstructed points and RMS point cloud alignment errors are shown in Tab. 11. Because Huang [2] has few points, the point cloud merge failed and is excluded from the table.

REFERENCES

- [1] P. J. Besl and N. D. McKay. Method for registration of 3-d shapes. In *Sensor Fusion IV: Control Paradigms and Data Structures*, vol. 1611, pp. 586–606. SPIE, 1992.
- [2] B. Huang, Y. Tang, S. Ozdemir, and H. Ling. A fast and flexible projector-camera calibration system. *IEEE Transactions on Automation Science and Engineering*, 18(3):1049–1063, 2020.
- [3] D. Moreno and G. Taubin. Simple, accurate, and robust projector-camera calibration. In *International Conference on 3d Imaging, Modeling, Processing, Visualization & Transmission*, pp. 464–471. IEEE, 2012.
- [4] N. Otsu. A threshold selection method from gray-level histograms. *IEEE Transactions on Systems, Man, and Cybernetics*, 9(1):62–66, 1979.
- [5] X. Xu, S. Xu, L. Jin, and E. Song. Characteristic analysis of otsu threshold and its applications. *Pattern Recognition Letters*, 32(7):956–961, 2011.

*e-mail: dongxin12345@email.swu.edu.cn

[†]e-mail: hling@cs.stonybrook.edu

[‡]e-mail: bhuang@swu.edu.cn. Corresponding author

Table 1: Comparison of color detection accuracy, #extracted nodes and decoded nodes among Huang's [2], Ours w/o MAP and Ours in different imaging settings.

Setting	Accuracy↑				#Extr. nodes↑				#Deco. nodes↑		
	Huang	Ours w/o MAP	Ours	Huang	Ours w/o MAP	Ours	Huang	Ours w/o MAP	Ours		
Setting1	0.9438	0.9635	0.9630	3388	3381	3381	2255	1992	1992		
Setting2	0.8837	0.9371	0.9522	3109	3102	3102	1428	1883	1889		
Setting3	0.8774	0.9535	0.9546	3578	3456	3456	1470	2104	2108		
Setting4	0.8353	0.9516	0.9503	3814	5010	5010	1064	2609	2609		
Setting5	0.9265	0.6996	0.9209	6788	8782	8782	3099	199	4424		
Setting6	0.9054	0.6420	0.8563	9938	10254	10254	4477	383	5147		
Setting7	0.7179	0.5947	0.8882	3916	4118	4118	771	466	2043		

Table 2: Comparison of color detection accuracy, #extracted nodes and decoded nodes among Huang's [2], Ours w/o MAP and Ours under different ambient light conditions.

Light	Accuracy↑				#Extr. nodes↑				#Deco. nodes↑		
	Huang	Ours w/o MAP	Ours	Huang	Ours w/o MAP	Ours	Huang	Ours w/o MAP	Ours		
Light1	0.7739	0.7221	0.9230	5498	5512	5512	1938	1016	3170		
Light2	0.7905	0.6998	0.8863	5487	5505	5505	1701	686	2950		
Light3	0.7938	0.6344	0.8462	5563	5571	5571	1993	571	2923		
Light4	0.7842	0.7056	0.8976	5699	5674	5674	1327	731	3034		
Light5	0.7198	0.6120	0.8857	7021	6993	6993	1077	248	3596		
Light6	0.7752	0.6494	0.8408	8290	8888	8888	2526	283	4549		

Table 3: Comparison of calibration reprojection errors among Moreno & Taubin [3], Huang [2], Ours w/o MAP and Ours in different imaging settings. Note, MT stands for Moreno & Taubin [3] and '—' stands for the calibration failed due to failed optimization.

Setting	RMSE ^c ↓						RMSE ^p ↓						RMSE ^{stereo} ↓		
	MT	Huang	Ours w/o MAP	Ours	MT	Huang	Ours w/o MAP	Ours	MT	Huang	Ours w/o MAP	Ours			
Setting1	0.1542	0.2403	0.2373	0.2373	0.6966	0.2614	0.2608	0.2608	0.5045	0.2511	0.2493	0.2493			
Setting2	0.1397	0.2165	0.2171	0.2213	1.2378	0.2201	0.2028	0.2049	0.8808	0.2183	0.2101	0.2132			
Setting3	0.2251	0.2428	0.2317	0.2322	0.8638	0.2271	0.2126	0.2129	0.6312	0.2351	0.2224	0.2228			
Setting4	0.1712	0.2619	0.2348	0.2348	2.6071	0.1790	0.1615	0.1615	1.8475	0.2243	0.2015	0.2015			
Setting5	0.4047	0.2487	—	0.2309	2.0323	0.1835	—	0.1680	1.4653	0.2185	—	0.2019			
Setting6	0.4904	0.3387	44.8472	0.2378	2.0266	0.5322	117.6157	0.2318	1.4744	0.4461	89.0077	0.2348			
Setting7	0.1290	12.4804	15.1692	0.2495	1.2099	13.5481	20.2279	0.2222	0.8603	13.0252	17.8784	0.2363			

Table 4: Comparison of calibration reprojection errors among Moreno & Taubin [3], Huang [2], Ours w/o MAP and Ours under different ambient light conditions. Note, MT stands for Moreno & Taubin [3] and '—' stands for the calibration failed due to failed optimization.

Light	RMSE ^c ↓						RMSE ^p ↓						RMSE ^{stereo} ↓		
	MT	Huang	Ours w/o MAP	Ours	MT	Huang	Ours w/o MAP	Ours	MT	Huang	Ours w/o MAP	Ours			
Light1	0.1639	0.2381	33.2597	0.2349	2.0851	0.1880	76.1839	0.1844	1.4790	0.2145	58.7801	0.2111			
Light2	0.1545	0.2352	29.0905	0.2353	1.9476	0.1884	82.7199	0.1883	1.3815	0.2131	62.0034	0.2131			
Light3	0.1764	0.2573	31.4283	0.2417	1.9410	0.2223	137.0736	0.1886	1.3781	0.2405	99.4407	0.2168			
Light4	0.1732	0.2291	101.4924	0.2243	2.1967	0.1920	64726.2026	0.1775	1.5582	0.2114	45768.3930	0.2023			
Light5	0.2181	1.7906	44.2268	0.2371	1.0249	83.9224	103.3141	0.1994	0.7409	59.3556	79.4664	0.2191			
Light6	0.1554	6.5740	—	0.2176	3.6754	7.6561	—	0.1761	2.6012	7.1356	—	0.1979			

Table 5: Comparison of point cloud alignment errors (mm) of David (mm) for Moreno & Taubin [3], Huang [2], Ours w/o MAP and Ours in different imaging settings. Note, MT stands for Moreno & Taubin [3] and '—' stands for the iterative closest point (ICP) [1] failed or reconstruction failed due to failed calibration.

Setting	Mean						Median						Standard deviation		
	MT	Huang	Ours w/o MAP	Ours	MT	Huang	Ours w/o MAP	Ours	MT	Huang	Ours w/o MAP	Ours			
Setting1	1.6517	1.5494	1.5404	1.5404	1.3724	1.3083	1.2983	1.2983	1.1786	1.0324	1.0278	1.0278			
Setting2	2.0382	1.5818	1.4402	1.4403	1.6409	1.2971	1.1713	1.1706	1.5942	1.1160	1.0192	1.0226			
Setting3	2.3749	1.6867	1.5806	1.5802	1.8226	1.4040	1.3378	1.3395	2.0222	1.1699	1.0498	1.0509			
Setting4	3.0323	1.6874	1.2942	1.2942	2.6606	1.2952	1.0101	1.0101	2.1332	1.3245	0.9872	0.9872			
Setting5	3.9359	1.5844	—	1.4829	3.3976	1.2710	—	1.1891	2.7709	1.1688	—	1.0826			
Setting6	2.9209	1.6296	—	1.5969	2.4442	1.2877	—	1.2674	2.1369	1.2040	—	1.1763			
Setting7	2.0357	13.1628	5.4138	1.6916	1.6413	10.8821	4.7273	1.4161	1.5116	10.9154	4.1503	1.1792			

Table 6: Comparison of point cloud alignment errors (mm) of girl (mm) for Moreno & Taubin [3], Huang [2], Ours w/o MAP and Ours in different imaging settings. Note, MT stands for Moreno & Taubin [3] and '—' stands for the iterative closest point (ICP) [1] failed or reconstruction failed due to failed calibration.

Setting	Mean				Median				Standard deviation			
	MT	Huang	Ours w/o MAP	Ours	MT	Huang	Ours w/o MAP	Ours	MT	Huang	Ours w/o MAP	Ours
Setting1	1.8939	1.7343	1.7249	1.7249	1.5983	1.4761	1.4694	1.4694	1.2860	1.1532	1.1430	1.1430
Setting2	1.9852	1.6296	1.3889	1.3885	1.7065	1.2782	1.0920	1.0953	1.3591	1.2391	1.0769	1.0731
Setting3	2.0830	1.6476	1.5968	1.5940	1.7342	1.3653	1.3344	1.3329	1.4713	1.1237	1.0812	1.0801
Setting4	4.1105	1.4405	1.1552	1.1552	3.9930	1.1436	0.9477	0.9477	2.5090	1.0319	0.7712	0.7712
Setting5	3.4947	1.7000	—	1.6081	2.7953	1.4089	—	1.3379	2.7879	1.1746	—	1.1013
Setting6	2.6449	1.9418	—	1.8981	2.1459	1.5997	—	1.5641	2.0528	1.4746	—	1.4289
Setting7	1.5500	15.9355	6.6280	1.5157	1.3049	12.8186	4.9851	1.2762	1.0187	12.3660	6.3522	0.9922

Table 7: Comparison of point cloud alignment errors (mm) of box (mm) for Moreno & Taubin [3], Huang [2], Ours w/o MAP and Ours in different imaging settings. Note, MT stands for Moreno & Taubin [3] and '—' stands for the iterative closest point (ICP) [1] failed or reconstruction failed due to failed calibration.

Setting	Mean				Median				Standard deviation			
	MT	Huang	Ours w/o MAP	Ours	MT	Huang	Ours w/o MAP	Ours	MT	Huang	Ours w/o MAP	Ours
Setting1	1.2292	1.1846	1.1785	1.1785	1.0422	1.0289	1.0203	1.0203	0.7817	0.7099	0.7077	0.7077
Setting2	1.2147	0.6707	0.6871	0.6759	1.0839	0.6168	0.6198	0.6135	0.6882	0.3371	0.3635	0.3526
Setting3	1.6082	0.8201	0.7697	0.7693	1.2960	0.7262	0.6979	0.6977	1.1212	0.4546	0.4003	0.4004
Setting4	2.8470	0.7664	0.7058	0.7058	2.2536	0.6519	0.6291	0.6291	2.1430	0.4779	0.3858	0.3858
Setting5	3.4014	0.9937	—	0.9227	2.9549	0.8623	—	0.8086	2.2771	0.5783	—	0.5195
Setting6	1.7350	0.9034	—	0.9150	1.4350	0.8042	—	0.8130	1.1942	0.4841	—	0.4957
Setting7	0.8195	15.3093	6.9558	0.7626	0.7312	10.8761	6.2259	0.7066	0.4497	13.9267	4.8298	0.3756

Table 8: Comparison of point cloud alignment errors (mm) of David (mm) for Moreno & Taubin [3], Huang [2], Ours w/o MAP and Ours under different ambient light conditions. Note, MT stands for Moreno & Taubin [3] and '—' stands for the reconstruction failed due to failed calibration.

Light	Mean				Median				Standard deviation			
	MT	Huang	Ours w/o MAP	Ours	MT	Huang	Ours w/o MAP	Ours	MT	Huang	Ours w/o MAP	Ours
Light1	2.8537	1.7245	19.9866	1.6993	2.3395	1.4508	16.3468	1.4012	2.1911	1.1837	15.9126	1.1997
Light2	2.5882	1.7112	18.8148	1.6949	2.1269	1.4088	12.8129	1.4036	1.9622	1.2079	18.9484	1.1792
Light3	2.5606	1.7474	41.5068	1.6879	2.1045	1.4695	36.0995	1.3936	1.9397	1.1959	28.8163	1.1867
Light4	1.8423	1.6323	23.1846	1.5484	1.4937	1.2924	18.0835	1.2317	1.4548	1.2113	19.3719	1.1362
Light5	2.4852	9.2549	22.4224	1.6698	2.0388	7.7165	18.7523	1.3862	1.9099	7.3835	16.7340	1.1737
Light6	2.3856	24.0016	—	1.4273	1.9974	20.0156	—	1.1259	1.7649	18.4220	—	1.0493

Table 9: Comparison of point cloud alignment errors (mm) of girl (mm) for Moreno & Taubin [3], Huang [2], Ours w/o MAP and Ours under different ambient light conditions. Note, MT stands for Moreno & Taubin [3] and '—' stands for the reconstruction failed due to failed calibration.

Light	Mean				Median				Standard deviation			
	MT	Huang	Ours w/o MAP	Ours	MT	Huang	Ours w/o MAP	Ours	MT	Huang	Ours w/o MAP	Ours
Light1	2.7717	1.5352	18.4845	1.5339	2.4223	1.2399	15.3412	1.2325	1.9680	1.0974	14.9082	1.1027
Light2	2.4978	1.4904	10.0544	1.5625	2.1560	1.2026	7.6564	1.2495	1.7804	1.0641	8.8892	1.1372
Light3	2.4932	1.5210	28.3107	1.5685	2.1467	1.2180	21.4328	1.2486	1.7762	1.0966	29.5077	1.1443
Light4	1.7747	1.3671	10.8638	1.3648	1.5069	1.1129	8.8445	1.1125	1.2078	0.9614	8.7018	0.9583
Light5	3.0365	11.8014	14.1339	1.5965	2.6889	8.4410	11.3291	1.3026	2.0501	11.3126	11.4128	1.1286
Light6	2.2320	12.7706	—	1.4002	1.8849	8.2654	—	1.1010	1.6335	17.7579	—	1.0309

Table 10: Comparison of point cloud alignment errors (mm) of box (mm) for Moreno & Taubin [3], Huang [2], Ours w/o MAP and Ours under different ambient light conditions. Note, MT stands for Moreno & Taubin [3] and '—' stands for the reconstruction failed due to failed calibration.

Light	Mean				Median				Standard deviation			
	MT	Huang	Ours w/o MAP	Ours	MT	Huang	Ours w/o MAP	Ours	MT	Huang	Ours w/o MAP	Ours
Light1	2.0242	1.0994	7.8760	0.9879	1.7807	0.9466	6.8096	0.8643	1.2878	0.6523	6.0891	0.5447
Light2	1.8009	0.9911	4.0474	1.0166	1.5790	0.8655	3.3318	0.8825	1.1242	0.5538	3.0599	0.5742
Light3	1.7836	1.0317	22.0487	1.0038	1.5616	0.8936	18.8423	0.8762	1.1136	0.5902	17.0161	0.5611
Light4	1.1591	0.9322	23.4383	0.8971	1.0171	0.8011	20.1127	0.7776	0.6627	0.5572	16.4948	0.5276
Light5	2.0851	19.2273	5.2254	0.8622	1.8998	13.2914	4.7108	0.7709	1.2650	17.4184	3.7681	0.4599
Light6	1.6624	6.7214	—	0.7413	1.3911	5.1888	—	0.6662	1.0883	5.3716	—	0.3904

Table 11: Comparison of the number of reconstructed points and RMS point cloud alignment errors (mm) among Huang [2], Ours w/o MAP, Ours w/o MAP (merged), Ours and Ours (merged), in a **carefully tuned** imaging setting and an **extreme** imaging setting.

Method	David (mm)				Fan (mm)				Box (mm)			
	#Points↑	Mean↓	Median↓	Std.↓	#Points↑	Mean↓	Median↓	Std.↓	#Points↑	Mean↓	Median↓	Std.↓
A carefully tuned setting												
Huang	500	1.22	0.97	0.80	310	1.04	0.83	0.73	192	0.68	0.61	0.46
Ours w/o MAP	1089	1.25	1.00	0.85	729	1.07	0.88	0.71	227	27.12	16.85	60.30
Ours w/o MAP (merged)	2708	2.09	1.46	1.94	1285	1.00	0.78	0.72	473	1.53	0.91	1.61
Ours	1113	1.26	1.00	0.87	729	1.07	0.88	0.71	704	0.76	0.65	0.70
Ours (merged)	2754	2.18	1.53	1.99	1314	1.00	0.81	0.70	1043	0.72	0.62	0.45
An extreme setting												
Huang	147	14.87	13.44	10.36	25	647.31	354.05	1038.08	91	22.85	19.17	16.58
Ours w/o MAP	733	2.18	1.32	19.03	147	73.29	44.76	102.88	110	60.12	24.54	127.02
Ours w/o MAP (merged)	2236	2.80	1.89	2.73	1274	1.70	1.26	1.51	715	1.47	0.82	4.46
Ours	887	1.48	1.35	0.94	689	1.39	1.22	0.84	559	0.92	0.78	0.55
Ours (merged)	2630	2.54	1.79	2.44	1437	1.31	1.00	1.02	1064	0.88	0.74	0.56